



DOSING FOR HEALTHCARE PROFESSIONALS

Use this chart to determine the proper dose of **TYLENOL**® for your patient. If possible, use weight to dose; otherwise, use age.[†]

[†] For pediatric products only

Recommended dose of **TYLENOL**® from your healthcare professional:

_____ dose of _____ product every _____ hours for _____ days

Visit Tylenol.com for additional product information

mL = milliliter

		Dosage							
		Infants' TYLENOL® Oral Suspension*	Children's TYLENOL® Oral Suspension*	Children's TYLENOL® Chewables*	Children's TYLENOL® Dissolve Packs	TYLENOL® Regular Strength Tablets & Liquid Gels	TYLENOL® Extra Strength Caplets, Coated Tablets, Rapid Release Gels, & Dissolve Packs	TYLENOL® For Children + Adults Family*	
Weight (lbs)	Age	Active ingredient: acetaminophen 160 mg (in each 5 mL)	Active ingredient: acetaminophen 160 mg (in each 5 mL)	Active ingredient: acetaminophen 160 mg (in each chewable tablet)	Active ingredient: acetaminophen 160 mg (in each powder)	Active ingredient: acetaminophen 325 mg (in each tablet/liquid gel)	Active ingredient: acetaminophen 500 mg (in each caplet /tablet /powder/gelcap)	Active ingredient: acetaminophen 160 mg (in each 5 mL)	
6-11 lbs	0-3 mos	1.25 mL	The markings on the included dosing cup correspond to the doses listed below. This cup is designed for dosing children ages 2 - 11 years.	-	Do not use	OTC label does not include dosing for children under 6 years old	OTC label does not include dosing for children under 12 years old	The markings on the included dosing cup correspond to the doses listed below. This cup is designed for dosing ages 2 years and older.	
12-17 lbs	4-11 mos	2.5 mL		-					
18-23 lbs	12-23 mos	3.75 mL		-					
24-35 lbs	2-3 yrs	5 mL	5 mL	1 tablet				5 mL	
36-47 lbs	4-5 yrs	The markings on the included syringe correspond to the doses listed above. This syringe is designed for dosing children ages 0 - 3 yrs.	7.5 mL	1½ tablets				7.5 mL	
48-59 lbs	6-8 yrs		10 mL	2 tablets	2 powders	Children 6 years to under 12 years: Take 1 tablet/liquid gel every 4 to 6 hours while symptoms last.		10 mL	Repeat dose every 4 hours while symptoms last. Do NOT give more than 5 doses in 24 hours.
60-71 lbs	9-10 yrs		12.5 mL	2½ tablets	2 powders	Do NOT take more than 5 liquid tablet/liquid gels in 24 hours		12.5 mL	
72-95 lbs	11 yrs		15 mL	3 tablets	3 powders			15 mL	
	Adults and children 12 years and over	-	-	-	-	Take 2 tablet/liquid gels every 4 to 6 hours while symptoms last. Do NOT take more than 10 tablet/liquid gels in 24 hours	Take 2 caplets/tablets/gelcaps/powders every 6 hours while symptoms last. Do NOT take more than 6 caplets/ tablets/gelcaps/ powders in 24 hours	20 mL	

* The OTC label says 'ask a doctor' for dosing of children under 2 years. All Infants' TYLENOL® and Children's TYLENOL® products have the same strength of acetaminophen: 160 mg (in each 5 mL, tablet, or powder).

Remind parents & caregivers:

- Read & follow the label on all **TYLENOL**® products.
- Repeat dose every 4 hours while symptoms last.
- Do NOT give more than 5 doses in 24 hours.
- Do NOT use with any other product containing acetaminophen.
- Use only the dosing device (syringe or dosing cup) that came with the product unless directed by a healthcare provider.

Remind patients:

- Read & follow the label on all **TYLENOL**® products.
- Do NOT use with any other product containing acetaminophen.
- Wait the right amount of time between doses